



## King's Research Portal

DOI:

[10.1016/j.scr.2016.01.004](https://doi.org/10.1016/j.scr.2016.01.004)

*Document Version*

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Miere, C., Hewitson, H., Devito, L., Wood, V., Kadeva, N., Cornwell, G., Codognotto, S., Stephenson, E., & Ilic, D. (2016). Generation of KCL018 research grade human embryonic stem cell line carrying a mutation in the DMPK gene. *Stem Cell Research*, 16(2), 342-344. <https://doi.org/10.1016/j.scr.2016.01.004>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.


### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



Lab Resource: Stem Cell Line

## Generation of KCL018 research grade human embryonic stem cell line carrying a mutation in the *DMPK* gene



Cristian Miere, Heema Hewitson, Liani Devito, Victoria Wood, Neli Kadeva, Glenda Cornwell, Stefano Codognotto, Emma Stephenson, Dusko Ilic\*

Stem Cell Laboratories, Division of Women's Health, Faculty of Life Sciences and Medicine, King's College London and Assisted Conception Unit, Guys' Hospital, London, United Kingdom

### ARTICLE INFO

#### Article history:

Received 30 December 2015

Received in revised form 5 January 2016

Accepted 12 January 2016

Available online 15 January 2016

### ABSTRACT


The KCL018 human embryonic stem cell line was derived from an embryo donated for research that carried an autosomal dominant mutation affecting one allele of the *DMPK* gene encoding the dystrophin myotonia protein kinase (2200 trinucleotide repeats; 14 for the normal allele). The ICM was isolated using laser microsurgery and plated on  $\gamma$ -irradiated human foreskin fibroblasts. Both the derivation and cell line propagation were performed in an animal product-free environment. Pluripotent state and differentiation potential were confirmed by in vitro assays.

© 2016 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

| Resource table  |   | (continued)                     |   |
|---|---|---------------------------------|---|
| Name of stem cell line  | KCL018  | Information in public databases | doi: 10.1038/nprot.2012.080<br><a href="http://www.ncbi.nlm.nih.gov/pubmed/22722371">http://www.ncbi.nlm.nih.gov/pubmed/22722371</a><br>KCL018 is a National Institutes of Health (NIH) registered hESC line<br>NIH Registration Number: 0218<br>NIH Approval Number: NIHhESC-13-0218<br><a href="http://grants.nih.gov/stem_cells/registry/current.htm?id=658">http://grants.nih.gov/stem_cells/registry/current.htm?id=658</a><br>The hESC line KCL018 is derived under license from the UK Human Fertilisation and Embryology Authority (research license numbers: R0075 and R0133) and also has local ethical approval (UK National Health Service Research Ethics Committee Reference: 06/Q0702/90).<br>Informed consent was obtained from all subjects and the experiments conformed to the principles set out in the WMA Declaration of Helsinki and the NIH Belmont Report.<br>No financial inducements are offered for donation. |
| Institution   | King's College London, London UK  | Ethics                          |   |
| Derivation team   | Neli Kadeva, Victoria Wood, Glenda Cornwell, Stefano Codognotto, Emma Stephenson  | Resource details                |   |
| Contact person and email  | Dusko Ilic, email: <a href="mailto:dusko.ilic@kcl.ac.uk">dusko.ilic@kcl.ac.uk</a>   | Consent signed                  | Aug 12, 2009  |
| Type of resource  | Biological reagent: cell line   | Embryo thawed                   | Aug 23, 2009  |
| Sub-type  | Human pluripotent stem cell line  | UK Stem Cell Bank               | Sep 23, 2010  |
| Origin  | Human embryo  | Deposit Approval                | Reference: SCSC10-30  |
| Key marker expression   | Pluripotent stem cell markers: NANOG, OCT4, TRA-1-60, TRA-1-81, alkaline phosphatase (AP) activity  | Sex                             | Female 46, XX   |
| Authentication  | Identity and purity of line confirmed<br>1) Ilic, D., Stephenson, E., Wood, V., Jacquet, L., Stevenson, D., Petrova, A., Kadeva, N., Codognotto, S., Patel, H., Semple, M., Cornwell, G., Ogilvie, C., Braude, P., 2012. Derivation and feeder-free propagation of human embryonic stem cells under xeno-free conditions. <i>Cytotherapy</i> . 14 (1), 122–128.   | Grade                           | Research  |
| Link to related literature (direct URL links and full references) | doi: 10.3109/14653249.2011.623692<br><a href="http://www.ncbi.nlm.nih.gov/pubmed/22029654">http://www.ncbi.nlm.nih.gov/pubmed/22029654</a><br>2) Stephenson, E., Jacquet, L., Miere, C., Wood, V., Kadeva, N., Cornwell, G., Codognotto, S., Dajani, Y., Braude, P., Ilic, D., 2012. Derivation and propagation of human embryonic stem cell lines from frozen embryos in an animal product-free environment. <i>Nat. Protoc.</i> 7 (7), 1366–1381. | Disease status (Fig. 1)         | Mutation affecting one allele of the <i>DMPK</i> gene encoding dystrophin myotonia protein kinase (~2200 CTG repeats; 14 for the normal allele) associated with Myotonic dystrophy Type 1 (Ilic et al., 2012)   |
|   |   | Karyotype (aCGH)                | No copy number changes detected   |
|   |   | DNA fingerprint                 | Allele sizes (in bp) of 17 microsatellite markers   |

\* Corresponding author.

E-mail address: [dusko.ilic@kcl.ac.uk](mailto:dusko.ilic@kcl.ac.uk) (D. Ilic).



**Fig. 1.** Genetic pedigree tree. The couple undergoing IVF had only one embryo in this particular cycle. The embryo carried the mutation and was donated for research.

|  |  |
|--|--|
| differentiation in vitro (immunostaining) (Fig. 3) | (tubulin, $\beta$ 3 class III); Mesoderm: ACTA2 (actin, $\alpha$ 2, smooth muscle) (Ilic et al., 2012) |
| Sibling lines available                            | No   |

We generated KCL018 clinical grade hESC line following protocols, established previously (Ilic et al., 2012; Stephenson et al., 2012). The expression of the pluripotency markers was tested after freeze/thaw cycle (Ilic et al., 2012). Differentiation potential into three germ layers was verified in vitro (Ilic et al., 2012).

## Materials and methods


### Consenting process

We distribute Patient Information Sheet (PIS) and consent form to the in vitro fertilization (IVF) patients if they opted to donate to research embryos that were stored for 5 or 10 years. They mail signed consent back to us and that might be months after the PIS and consent were mailed to them. If in the meantime new versions of PIS/consent are implemented, we do not send these to the patients or ask them to resign; the whole process is done with the version that was given them initially. The PIS/consent documents (PGD-V.6) were created on Aug. 10, 2007. HFEA Code of Practice that was in effect at the time of document creation: Edition 7 – R.1 (<http://www.hfea.gov.uk/2999.html>). The donor couple signed the consent on Oct. 15, 2009. HFEA Code of Practice that was in effect at the time of donor signature: Edition 8 – R.1. HFEA Code of Practice Edition 7 – R.1 was in effect until Dec. 09, 2007 and Edition 8 – R.1 was in effect: Oct. 01, 2009–Apr. 06, 2010.


### Embryo culture and micromanipulation

Embryo culture and laser-assisted dissection of inner cell mass (ICM) were carried out as previously described in details (Ilic et al., 2012; Stephenson et al., 2012). The cellular area containing the ICM was then washed and transferred to plates containing mitotically inactivated human neonatal foreskin fibroblasts (HFF).

|   |  |
|---|--|
| Viability testing                             | specific for chromosomes 13, 18 and 21 (Ilic et al., 2012)       |
| Pluripotent markers (immunostaining) (Fig. 2) | Pass   |
| Three germ layers                             | NANOG, OCT4, TRA-1-60, TRA-1-81, AP activity (Ilic et al., 2012) |
|   | Endoderm: AFP ( $\alpha$ -fetoprotein); Ectoderm: TUBB3          |



**Fig. 2.** Expression of pluripotency markers. Pluripotency is confirmed by immunostaining (Oct4, Nanog, TRA-1-60, TRA-1-81) and alkaline phosphatase (AP) activity assay. Actin stress fibers, visualized with rhodamine-phalloidin (red), are present in both feeders and hES cell colonies, whereas AP activity (green) is detected only in hES cells. Scale bar, 50  $\mu$ m.



**Fig. 3.** Differentiation of three germ layers in vitro is confirmed by detection of markers: smooth muscle actin (red) for mesoderm,  $\beta$ -III tubulin (red) for ectoderm and  $\alpha$ -fetoprotein (red) for endoderm. Nuclei are visualized with Hoechst 33,342 (blue). Scale bar, 50  $\mu$ m.

### Cell culture

ICM plated on mitotically inactivated HFF were cultured as described (Ilic et al., 2012; Stephenson et al., 2012). TE cells were removed mechanically from outgrowth (Ilic et al., 2007; Ilic et al., 2010). hESC colonies were expanded and cryopreserved at the third passage.

### Viability test

Straws with the earliest frozen passage (p. 2–3) are thawed and new colonies are counted three days later. These colonies are then expanded up to passage 8, at which point cells were part frozen and part subjected to standard battery of tests (pluripotency markers, in vitro and in vivo differentiation capability, genetics, sterility, mycoplasma).

### Pluripotency markers

Pluripotency was assessed using two different techniques: enzymatic activity assay [alkaline phosphatase (AP) assay] and immunostaining as described (Ilic et al., 2012; Stephenson et al., 2012).

### Genotyping

DNA was extracted from hES cell cultures using a Chemagen DNA extraction robot according to the manufacturer's instructions. Amplification of polymorphic microsatellite markers was carried out as described (Ilic et al., 2012). Allele sizes were recorded to give a unique fingerprint of each cell line.

### Array comparative genomic hybridization (aCGH)

aCGH was performed as described in details (Ilic et al., 2012).

| Chr | Marker   | Allele 1 | Allele 2 |
|-----|----------|----------|----------|
| 13  | D13S252  | 299      | 299      |
|     | D13S305  | 443      | 458      |
|     | D13S325  | 289      | 297      |
|     | D13S628  | 429      | 450      |
|     | D13S634  | 397      | 417      |
|     | D18S386  | 383      | 386      |
|     | D18S390  | 372      | 372      |
| 18  | D18S391  | 209      | 217      |
|     | D18S535  | 482      | 486      |
|     | D18S819  | 408      | 424      |
|     | D18S976  | 479      | 479      |
|     | D18S978  | 219      | 219      |
|     | D21S11   | 244      | 251      |
|     | D21S1409 | 216      | 224      |
| 21  | D21S1411 | 308      | 308      |
|     | D21S1435 | 184      | 188      |
|     | D21S1437 | 331      | 335      |

### Author disclosure statement

There are no competing financial interests in this study.

### Acknowledgments

This work was supported by the UK Medical Research Council grants G0701172 and G0801061. We thank Dr. Yacoub Khalaf, Director of the Assisted Conception Unit of Guy's and St Thomas' NHS Foundation Trust and his staff for supporting the research program. We are especially indebted to Prof Peter Braude and to the patients who donated embryos.

### References

- Ilic, D., Stephenson, E., Wood, V., Jacquet, L., Stevenson, D., Petrova, A., Kadeva, N., Codognotto, S., Patel, H., Semple, M., Cornwell, G., Ogilvie, C., Braude, P., 2012. Derivation and feeder-free propagation of human embryonic stem cells under xeno-free conditions. *Cytotherapy* 14 (1), 122–128.
- Ilic, D., Caceres, E., Lu, S., Julian, P., Foulk, R., Krtolica, A., 2010. Effect of karyotype on successful human embryonic stem cell derivation. *Stem Cells Dev.* 19 (1), 39–46.
- Ilic, D., Genbacev, O., Krtolica, A., 2007. Derivation of hESC from intact blastocysts. *Curr. Protoc. Stem Cell Biol.* (Chapter 1: Unit 1 A.2).
- Stephenson, E., Jacquet, L., Miere, C., Wood, V., Kadeva, N., Cornwell, G., Codognotto, S., Dajani, Y., Braude, P., Ilic, D., 2012. Derivation and propagation of human embryonic stem cell lines from frozen embryos in an animal product-free environment. *Nat. Protoc.* 7 (7), 1366–1381.